

Eco-Bb helps control whitefly and suppresses red spider mite on beans, tomatoes, cucumbers and brinjals under greenhouse and field conditions. It is also registered for the control of Potato Weevil on Potatoes.

WARNINGS

- Keep out of reach of children, uninformed persons and animals.
- Store in a refrigerator away from chemical pesticides. (Do not freeze)
- May cause slight eye irritations.
- May cause allergic reactions in sensitive persons.

PRECAUTIONS

- **Remember you are working with a living organism.**
- Do not use after expiry date.
- Do not mix in containers that have previously been used

DIRECTIONS FOR USE

- Mix with water at a rate of 1g / L water and apply as a full cover spray every 7-14 days depending on severity of pest.
- Preferably apply product when humidity is high and UV is low - i.e. late afternoon.
- Good spray coverage is essential. Aerial applications are successful as long as good coverage is achieved.
- Wetters and other adjuvants are not needed unless recommended.
- Ensure areas where pests are habitually found are covered. i.e. undersides of leaf or in leaf folds.
- Avoid the use of broad spectrum fungicides for at least three days before and after Eco-Bb applications.
- Close the lid of the bottle firmly after the seal is broken

for toxic substances. Wash the containers well with soap and water before use.

- Use within 6 hours of mixing.
- Do not use around bee hives or where bees are actively foraging.
- Shake or stir the fungus mixture frequently during use to avoid settling out of the product.
- Do not leave powder or mixture standing in the sun.
- Do not mix fungus with any agrochemicals without first consulting your sales representative.
- Avoid inhaling dust.
- Wash hands with soap and water before eating, drinking or smoking
- Do not use or apply when temperatures are below 15°C or above 35°C

Suggested Application Rates and Frequency:

Objective	Recommended Rate	Frequency
Preventative Control	1g / L water (300 - 600g depending on spray volume, crop and growth stage)	Apply weekly as a full cover spray to foliage. Ensure areas where pests are habitually found are covered. i.e. underside of leaf or leaf folds.
Curative Control	1g / L water (300 - 900g depending on spray volume, crop and growth stage)	Initial application of 900g / Ha followed up at least twice at 600g / Ha at 3-5 day intervals or until pest is under adequate control. Maintain control with preventative program especially when environmental conditions are favourable to pest outbreak.
Preventative Program with AgriSil	1g / L water (min 300g / Ha Eco-Bb) Plus 1-2 L AgriSil K50 / Ha	Apply weekly, alternating applications of Eco-Bb and AgriSil. This program allows for the application of broad spectrum fungicides which can be applied in the same week as AgriSil without negatively affecting the Eco-Bb or compromising disease and pest control. AgriSil will also aid plant disease resistance.

Disclaimer: Although this inoculant has been tested under a variety of conditions, the supplier does not warrant that it will be efficacious under all conditions. Because the active agent is a living organism, its action may be affected by storage, handling, climatic or soil conditions. This naturally occurring organism should not cause damage to crops, vegetation, or the environment, nor harm to man and animals. However, in the unlikely event of this happening, the supplier does not accept responsibility for any damage that may occur.

Marketed and Distributed by:



Madumbi Sustainable Agriculture (Pty) Ltd
Postnet Suite 148, Private Bag X9118
Pietermaritzburg 3200
Tel. +27 (0) 86 104 5261
support@madumbi.co.za
www.madumbi.co.za

Manufactured by:

