

The fungus *Trichoderma harzianum* helps control root diseases and enhances the growth of a wide variety of vegetables, crops, ornamentals and Eucalyptus species.

### WARNINGS

- Keep out of reach of children, uninformed persons and animals.
- Store in a refrigerator away from chemical pesticides.
- May cause slight eye irritations.
- May cause allergic reactions in sensitive persons.

### PRECAUTIONS

- **Remember you are working with a living organism.**
- Do not use after expiry date.
- Do not mix in containers that have previously been used

### DIRECTIONS FOR USE

**Seed Treatment:** Seed dressing application rates differ for different crops. Refer to table below. A suitable sticker may be used if necessary and is available from the supplier. A sticker is not required where Eco-T is added during on-farm treatment with insecticides, compatible fungicides or *Rhizobium* inoculant. Apply as a dry powder to slightly moist seed.

**Drench:** Apply as a drench to seedling trays, seedbeds, potted plants and turf at planting and monthly thereafter. Mix at a rate of 1 g per 4 L water. Mix the required amount of powder to a thin paste with a small volume of water before adding the remaining water. Irrigate the fungus into the root zone after application.

- The contents of a 250 g bottle will treat approximately 3000 m<sup>2</sup> of seedbed or 3000 seedling trays.
- For severe soil borne disease problems in field crops use a field application of 500-750 g/Ha.
- For further information on application under specific circumstances or fungicide compatibility consult your agent.

### Suggested Application Rates and Frequency:

Crop	Seed dressing		Field Application Rate	Frequency
	Rate per kg seed	Rate per Ha		
Brassica	40 g	100-125 g	Drench 1 g per 4 L water	At plant or before transplant. Repeat monthly 250 g/Ha
Carrot	40 g		250 g/Ha	Repeat monthly
Cucumber	5 g		Drench 1 g per 4 L water	Repeat monthly 250 g/Ha
Dry Bean	1-2 g	80 g		Seed treatment before sowing
Flowers		250 g	Drench 1 g per 4 L water	Repeat monthly / bimonthly 250 g/Ha
Sorghum	5 g	100-120 g (Forage) 40-50 g (Grain)		Seed treatment before sowing
Hydroponics (Lettuce)	40 g		1 g per 8 L water for recycling	Weekly top-up
Lucerne	7.5 g	90 g		With <i>Rhizobium</i> before planting
Maize	1 g	25-50 g	80-100 g/Ha in furrow or 250 g/Ha via pivot	Just prior to planting
Orchard Crops			500 g 1st applic. & 250-500 g 2nd and 3rd	At least 3 x per season
Pasture	5 g			At planting
Peppers / Tomatoes	20 g		Drench seedling before transplant 1 g per 4 L water	Repeat monthly 250 g/Ha
Potatoes		500 g + 2 kg Kaolin (diluent)	250 g/Ha	At planting and 28 days after emergence.
Soybean	1 g	50-75 g		With <i>Rhizobium</i> before planting
Strawberries			500 g/Ha	Monthly
Wheat	1-2 g	40-80 g		Just prior to planting
Sunflower	5 g	15-20 g		Just prior to planting

**Disclaimer:** Although this inoculant has been tested under a variety of conditions, the supplier does not warrant that it will be efficacious under all conditions. Because the active agent is a living organism, its action may be affected by storage, handling, climatic or soil conditions. This naturally occurring organism should not cause damage to crops, vegetation, or the environment, nor harm to man and animals. However, in the unlikely event of this happening, the supplier does not accept responsibility for any damage that may occur.

Marketed and Distributed by:



Madumbi Sustainable Agriculture (Pty) Ltd  
Postnet Suite 148, Private Bag X9118  
Pietermaritzburg 3200  
Tel. +27 (0) 86 104 5261  
support@madumbi.co.za  
www.madumbi.co.za

Manufactured by:

