

A member of the Andermatt group

A *Bacillus thuringiensis* subspecies kurstaki (Btk) flowable granule concentrate formulation for the control of various Lepidoptera larvae on crops as indicated.

WARNINGS

- May irritate skin and eyes.
- Keep out of reach of children, uninformed persons and animals.
- Btk particles are sensitive to UV and prolonged exposure can result in reduced efficacy.
- Do not leave the product or mixture standing in the sun.
- Delfin WG will remain viable for at least 24 months stored at room temperature.

PRECAUTIONS

- Avoid direct contact with eyes. In case of accidental contact immediately flush eyes with clean water.

DIRECTIONS FOR USE

COMPATIBILITY

Delfin WG contains Btk particles and can be tank mixed with most pesticides and foliar feeds. A jar test of any combination is recommended prior to tank mixing to ensure physical compatibility. If you are uncertain please contact Madumbi Sustainable Agriculture.

MIXING INSTRUCTIONS

- Fill spray tank with half the required water volume and start agitator mechanism.
- Add the correct amount of product to the water in the tank at the recommended dose rate as indicated in the instructions.
- Add any other spray material if necessary followed by the balance of required water volume.
- Apply as a full cover film spray using conventional spray equipment. Good coverage is essential.

IMPORTANT POINTS TO REMEMBER

- Due to UV sensitivity it is recommended that Delfin is applied during the afternoon or evenings.
- Delfin WG is a wettable granule formulation and the Btk spores and crystals should be kept in suspension.
- Mixed volumes must be applied as soon as possible and not left overnight or standing in the sun.
- Delfin WG does not impose any risk of residues on harvested crops.
- Delfin WG will not harm any beneficial organisms
- Minimum water volume for ground application 150 L/Ha and for aerial application 25 L/Ha.

REGISTERED USAGES:

Crop / Pest	Dosage	Remarks
Lettuce African Bollworm (<i>Helicoverpa armigra</i>)	1 kg / ha	Application timing- and intervals <ul style="list-style-type: none"> • 1st DELFIN WG with first detection of specified pest eggs. • 2nd DELFIN WG application 7 -10 days after first application. • Continue application of DELFIN WG every 7 – 10 days as long as specified pest is present. Application guidelines <ul style="list-style-type: none"> • Apply through conventional application systems as described in the mixing instructions. DELFIN WG should be applied as a full cover film spray. Coverage of feeding areas essential.
Tomato, Pepper, Eggplant, Potatoe <i>Tuta absoluta</i>		
Cotton, Cruciferae, Maize, Sorghum, Soya, Wheat Fall armyworm (<i>Spodoptera frugiperda</i>)		

Disclaimer: Although this product has been tested under a variety of conditions, the supplier does not warrant that it will be efficacious under all conditions. Because the active agent is a living organism, its action may be affected by storage, handling, climatic or soil conditions. This naturally occurring organism should not cause damage to crops, vegetation, or the environment, nor harm to man and animals. However, in the unlikely event of this happening, the supplier does not accept responsibility for any damage that may occur.

Marketed and Distributed by:



Madumbi Sustainable Agriculture (Pty) Ltd
Postnet Suite 148, Private Bag X9118
Pietermaritzburg 3200
Tel. +27 (0) 86 104 5261
support@madumbi.co.za
www.madumbi.co.za

Manufactured by:



Backed by science, loved by nature

Delfin WG

Application Instructions

- Avoid contact with skin, wash hands with soap after application and before consuming food, drinking or smoking.
- Do not inhale spray mist.
- Do not eat, drink or smoke whilst mixing or applying.
- Prevent contamination of food, foodstuffs, drinking water and eating utensils.
- Clean applicator before and after use. Do not contaminate other crops, grazing, rivers, dams and areas not under treatment with wash water
- Store in original container in a dry area, away from food and feed stuffs.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsing water to the contents of the spray or mixing tank. Destroy the container, and do not use empty container for any other purpose.